

Gene-Specific Differential DNA Methylation and Chronic Arsenic Exposure in an Epigenome-Wide Association Study of Adults in Bangladesh

Maria Argos, Lin Chen, Farzana Jasmine, Lin Tong, Brandon L. Pierce, Shantanu Roy, Rachelle Paul-Brutus, Mary V. Gamble, Kristin N. Harper, Faruque Parvez, Mahfuzar Rahman, Muhammad Rakibuz-Zaman, Vesna Slavkovich, John A. Baron, Joseph H. Graziano, Muhammad G. Kibriya, and Habibul Ahsan

<http://dx.doi.org/10.1289/ehp.1307884>

Received: 13 November 2013

Accepted: 15 October 2014

Advance Publication: 17 October 2014

This article will be available in its final, 508-conformant form 2–4 months after Advance Publication. If you require assistance accessing this article before then, please contact [Dorothy L. Ritter](#), *EHP* Web Editor. *EHP* will provide an accessible version within 3 working days of request.

Gene-Specific Differential DNA Methylation and Chronic Arsenic Exposure in an Epigenome-Wide Association Study of Adults in Bangladesh

Maria Argos,¹ Lin Chen,¹ Farzana Jasmine,¹ Lin Tong,¹ Brandon L. Pierce,¹ Shantanu Roy,¹ Rachelle Paul-Brutus,¹ Mary V. Gamble,² Kristin N. Harper,² Faruque Parvez,² Mahfuzar Rahman,³ Muhammad Rakibuz-Zaman,³ Vesna Slavkovich,² John A. Baron,⁴ Joseph H. Graziano,² Muhammad G. Kibriya,¹ and Habibul Ahsan^{1,5}

¹Department of Health Studies, The University of Chicago, Chicago, Illinois, USA; ²Department of Environmental Health Sciences, Mailman School of Public Health, Columbia University, New York, New York, USA; ³U-Chicago Research Bangladesh (URB), Ltd., Dhaka, Bangladesh; ⁴Department of Medicine, University of North Carolina School of Medicine, Chapel Hill, North Carolina, USA; ⁵Departments of Medicine and Human Genetics and Comprehensive Cancer Center, The University of Chicago, Chicago, Illinois, USA

Address correspondence to Habibul Ahsan, The University of Chicago, Department of Health Studies, 5841 South Maryland Avenue, MC2007, Chicago, IL 60637 USA. Telephone: 773-834-9956. E-mail: habib@uchicago.edu

Running title: Arsenic and epigenome-wide DNA methylation

Acknowledgments: We thank Devin Koestler for assistance and helpful discussion regarding the cell type deconvolution analyses. This work was supported by the National Institutes of Health grant number R01 CA107431.

Competing financial interests: The authors declare they have no actual or potential competing financial interests.

Abstract

Background: Inorganic arsenic is one of the most common naturally occurring contaminants found in the environment. Arsenic is associated with a number of health outcomes, with epigenetic modification suggested as a potential mechanism of toxicity.

Objective: Among a sample of 400 adult participants, we evaluated the association between arsenic exposure, as measured by blood and urinary total arsenic concentrations, and epigenome-wide white blood cell DNA methylation.

Methods: Linear regression models were used to examine the associations between arsenic exposure and methylation at each CpG site, adjusted for sex, age, and batch. Differentially methylated loci were subsequently examined in relation to corresponding gene expression for functional evidence of gene regulation.

Results: In adjusted analyses, we observed 4 differentially methylated CpG sites with urinary total arsenic concentration and 3 differentially methylated CpG sites with blood arsenic concentration based on the Bonferroni-corrected significance threshold of $P < 1 \times 10^{-7}$. Methylation of *PLA2G2C* cg04605617 was the most significantly associated locus in relation to both urinary ($P = 3.40 \times 10^{-11}$) and blood arsenic concentrations ($P = 1.48 \times 10^{-11}$). Three additional novel methylation loci in *SQSTM1* (cg01225779), *SLC44A4* (cg06121226), and *IGH* (cg13651690) were also significantly associated with arsenic exposure. Further, there was evidence of methylation-related gene regulation based on gene expression for a subset of differentially methylated loci.

Conclusions: Significant associations between arsenic exposure and gene-specific differential white blood cell DNA methylation were observed in this study, suggesting that epigenetic

modifications may be an important pathway underlying arsenic toxicity. The specific differentially methylated loci identified may inform potential pathways for future interventions.

Introduction

Millions of individuals worldwide are exposed to inorganic arsenic through drinking water as well as dietary sources (Smith et al. 2000). Arsenic is a well-established human carcinogen (IARC Working Group on the Evaluation of Carcinogenic Risks to Humans 2012); however, the exact mechanism by which it causes cancer has not been established (Kitchin and Conolly 2010). There is both *in vitro* and *in vivo* evidence to suggest that epigenetic alterations may mediate arsenic toxicity, as recently reviewed in (Reichard and Puga 2010; Ren et al. 2011).

Several human studies have examined global DNA methylation in blood in relation to arsenic exposure using surrogate markers of global DNA methylation, such as long interspersed nucleotide element-1, Alu element methylation, methyl incorporation assays, or luminometric methylation assays. The findings from these studies have largely been inconsistent and included a number of differences in exposure measures and doses across studies (Hossain et al. 2012; Intarasunanont et al. 2012; Kile et al. 2012; Lambrou et al. 2012; Majumdar et al. 2010; Pilsner et al. 2007, 2009; Pilsner et al. 2012; Tajuddin et al. 2013; Wilhelm et al. 2010). A number of studies have evaluated arsenic in relation to gene-specific DNA methylation, most frequently assessing p16 and p53 promoter methylation (Chanda et al. 2006; Chen et al. 2007; Engstrom et al. 2013; Hossain et al. 2012; Intarasunanont et al. 2012; Marsit et al. 2006; Zhang et al. 2007). However, relatively few epigenome-wide DNA methylation studies have been conducted to investigate epigenetic alterations of arsenic toxicity in humans, evaluating associations with arsenical skin lesion status (Seow et al. 2014; Smeester et al. 2011), urinary arsenic species (Bailey et al. 2013), *in utero* arsenic exposure (Kile et al. 2014; Koestler et al. 2013a), toenail arsenic concentration (Liu et al. 2014), and arsenic-related urothelial carcinomas (Yang et al.

2014). Smeester et al. (Smeester et al. 2011) examined epigenome-wide promoter DNA methylation in peripheral blood leukocytes among 16 arsenic-exposed females from Mexico in relation to skin lesion status, and observed 183 differentially methylated genes, of which 182 were hypermethylated. Bailey et al. (Bailey et al. 2013) evaluated these data in relation to urinary arsenic species and observed nominally significant differential promoter DNA methylation in 812 unique genes, of which the majority were hypomethylated, in relation to relative urinary arsenic metabolite species. Koestler et al. (Koestler et al. 2013a) evaluated *in utero* arsenic exposure in relation to epigenome-wide cord blood methylation in 134 US-based individuals and observed evidence of enrichment of hypermethylated loci in CpG islands as well as a suggested indication of endocrine disruption effects of arsenic through hypomethylation of *ESR1* and *PPARGC1A*.

This previous evidence warrants further investigation of arsenic exposure on gene-specific DNA methylation in a comprehensive manner and in a larger study population for potential identification of mechanisms associated with arsenic-related toxicity. We conducted an epigenome-wide association study among 400 Bangladeshi individuals with manifest arsenical skin lesions to assess whether arsenic exposure level (as measured by blood arsenic and urinary total arsenic concentrations) is associated with differential white blood cell DNA methylation.

Methods

Study population

The Bangladesh Vitamin E and Selenium Trial is a 2×2 factorial randomized chemoprevention trial evaluating the long-term effects of vitamin E and selenium supplementation on non-melanoma skin cancer risk (Argos et al. 2013). Participants are residents of rural communities in

central Bangladesh. Eligibility criteria included age between 25 and 65 years, permanent residence in the study area, manifest arsenical skin lesions, and no prior cancer history. Between April 2006 and August 2009, a total of 7,000 individuals were enrolled into the study. Trained study physicians, blinded to participants' arsenic exposure, conducted in-person interviews and clinical evaluations, and collected urine and blood samples from participants in their homes using structured protocols. There were 413 participants randomly sampled for epigenome-wide methylation analyses, on whom baseline biological specimens collected prior to the start of the trial intervention were utilized.

The study protocol was approved by the relevant Institutional Review Boards in the US (The University of Chicago and Columbia University), and Bangladesh (Bangladesh Medical Research Council). Informed consent was provided by participants prior to the baseline interview of the original study.

Exposure assessment

Urinary total arsenic concentration was measured in the baseline spot urine sample by graphite furnace atomic absorption spectrometry (Analyst 600, Perkin Elmer, Norwalk, CT, USA) with a detection limit of 2 µg/L, in a single laboratory (Trace Metal Core Laboratory at Columbia University) (Nixon et al. 1991). Urinary creatinine was also measured for all participants in the same laboratory by a colorimetric method based on the Jaffe reaction (Heinegard and Tiderstrom 1973). Urinary total arsenic was divided by creatinine to obtain a creatinine-adjusted urinary total arsenic concentration, expressed as µg/g creatinine. Creatinine-adjusted urinary total arsenic, a good biomarker of aggregate ingested arsenic exposure, captures exposure from all sources including water, food, soil, and dust (Hughes 2006).

Venous whole blood samples collected at baseline were analyzed for blood arsenic concentration using a Perkin Elmer Elan DRC (Dynamic Reaction Cell) II ICP-MS equipped with an AS 93+ autosampler (Perkin Elmer, Norwalk, CT, USA). ICP-MS-DRC methods for metals in whole blood were developed according to published procedures (Pruszkowski et al. 1998; Stroh 1988), with modifications for blood sample preparation as suggested by the Laboratory for ICP-MS Comparison Program (Institut National de Sante Publique du Québec).

DNA methylation

DNA was extracted by DNeasy Blood Kits (Qiagen, Valencia, CA, USA). Bisulfite conversion was performed using the EZ DNA Methylation Kit (Zymo Research, Irvine, CA, USA). DNA methylation was measured using the Illumina HumanMethylation 450K BeadChip using 500 ng of bisulfite-converted DNA per sample according to the manufacturer's protocol, which interrogates 485,577 CpG sites. The methylation score for each CpG site was represented as the β value, on a continuous scale between 0 (unmethylated) to 1 (completely methylated), and was quantile normalized. Among the 413 participants for whom DNA methylation data was generated, we excluded 6 samples whose reported sex did not correspond with predicted sex based on methylation patterns of the X and Y chromosomes and 7 samples with >5% of CpGs containing missing values or P for detection >0.05. This resulted in 400 samples retained for analyses. We removed β values that were associated with a P for detection value >0.05. We also excluded probes on the X (n=11,232) and Y (n=416) chromosomes, probes with missing chromosome data (mostly control probes) (n=65), and probes with >10% missing data across samples (n=1,932). This resulted in 471,932 probes included in the statistical analyses. Quantile normalized β values were logit transformed and adjusted for batch variability using ComBat

software (Johnson et al. 2007). Based on 11 samples run in duplicate across two different plates in these experiments, the average inter-assay Spearman correlation coefficient was 0.987 (range: 0.974–0.993).

Gene expression

RNA was extracted from mononuclear cells preserved in RLT buffer, stored at -80°C, using RNeasy Micro Kit from Qiagen (Valencia, CA, USA). The concentration and quality of RNA was checked on Nanodrop 1000. cRNA synthesis was done from 250 ng of RNA using Illumina TotalPrep 96 RNA Amplification kit. Gene expression was measured using the Illumina HumanHT-12-v4 BeadChip utilizing 750 ng of cRNA according to the manufacturer's protocol. The chip contains a total of 47,231 probes covering 31,335 genes. Quantile normalized expression values were log₂ transformed and adjusted for batch variability using ComBat software (Johnson et al. 2007). Gene expression data were available for the 400 individuals included in these analyses.

Genotyping

Genotyping procedures have been described previously in detail (Pierce et al. 2012). Briefly, DNA extraction was carried out from whole blood using the QIAamp 96 DNA Blood Kit from Qiagen (Valencia, CA, USA). Any DNA sample with a concentration <40 ng/μL, and/or 260/280 ratio outside the range of <1.6 to ≥2.1 (measured by Nanodrop 1000), and/or fragmented DNA <2 Kb (assessed by smearing in Agilent BioAnalyzer) was excluded. Genotyping was performed using the Illumina HumanCytoSNP-12 BeadChip utilizing 250 ng DNA according to the manufacturer's protocol. Using 257,768 genotyped SNPs after quality control (QC) procedures, we performed imputation using MaCH on the basis of the HapMap 3 Gujarati Indians in Houston

(GIH) population (Build 36), yielding 1,211,988 million SNPs after QC procedures. Genotype data were available for 393 individuals included in these analyses.

Replication study

For replication of our top differentially methylated loci, we report associations from an independent sample of 48 Bangladeshi adult males from the ongoing Folate and Creatinine Trial (FACT), prior to intervention. No participants in this sample had manifest arsenical skin lesions. A detailed description of the methods used to measure and analyze DNA methylation in this study can be found in Harper et al. (Harper et al. 2013). Briefly, water arsenic concentration was measured for all study participants at Columbia University as previously described (Van Geen et al. 2002), and individuals were categorized as low (50–100 $\mu\text{g/L}$, $n=25$) or high exposure (> 100 $\mu\text{g/L}$, $n=23$) for statistical analyses. After Ficoll separation, DNA from peripheral blood mononuclear cells (PBMCs) was extracted using the 5 PRIME ArchivePure DNA Blood Kit (Fisher Scientific, Pittsburgh, PA, USA). Epigenome-wide methylation of PBMC DNA was measured using the Illumina HumanMethylation 450K BeadChip (Roswell Park Cancer Institute Laboratory). Data were processed using a standard quality control protocol and adjusted for batch effects prior to analyses using COMBAT (Johnson et al. 2007). Associations from linear regression models are reported.

Statistical analyses

For each CpG site, a separate linear regression model was run regressing the logit-transformed β value on continuous arsenic exposure (i.e., blood arsenic or urinary total arsenic concentration), sex, and age. We present model coefficients and standard errors from the linear regression models comparing the 75th versus 25th percentile of each arsenic distribution. To correct for

multiple comparisons, we considered a Bonferroni corrected ($P < 1 \times 10^{-7}$) significance threshold, however present results for all loci with $P < 1 \times 10^{-5}$. For differentially methylated probes with $P < 1 \times 10^{-5}$, we used linear regression to examine the association of methylation with corresponding RNA transcript levels of the gene containing the methylation locus, adjusted for sex, age, and urinary total arsenic concentration. Methylation and expression data were adjusted for batch effects prior to analyses using COMBAT (Johnson et al. 2007).

COMBAT batch-adjusted methylation data were used to infer white blood cell type fractions using the statistical method of Houseman et al. (Houseman et al. 2012). Briefly, we utilized 466 methylation probes that had been previously identified to be associated with leukocyte distributions (Houseman et al. 2012) to infer the distribution of white blood cell types in our study samples. Linear regression was used to evaluate the association between quartiles of arsenic exposure, based on the distribution in our study sample, in relation to the estimated cell type fractions adjusted for sex and age.

To evaluate previously reported associations from other studies, we conducted a lookup of individual CpG loci of interest within our dataset. For these analyses a CpG locus associated with $P < 0.05$ was considered to be statistically significant. The R program MethLAB v1.5 (Kilaru et al. 2012) and SAS (Cary, NC, USA) were used to run all analyses.

Results

Study sample characteristics are summarized in Table 1. For the present analysis, we ran models separately for blood and urinary total arsenic concentrations (Spearman correlation coefficient=0.91). We identified 3 loci that were significantly differentially methylated in

relation to blood arsenic concentration based on the Bonferroni threshold $P < 1 \times 10^{-7}$, as shown in Figure 1. We identified 4 loci that were significantly differentially methylated in relation to urinary total arsenic concentration based on the Bonferroni threshold $P < 1 \times 10^{-7}$, as shown in Figure 2. The loci based on the Bonferroni threshold were common between the two analyses, and several overlapping associations that did not meet the strict Bonferroni threshold, but had $P < 1 \times 10^{-5}$ were also observed between the exposure analyses, as summarized in Table 2. Furthermore, there appeared to be enrichment for differentially methylated loci in relation to urinary and blood arsenic concentrations in ocean (isolated CpG loci in the genome) and CpG island shore regions (within 2kb from a CpG island; Supplemental Material, Figure S1). Only a single gene contained more than one differentially methylated locus with $P < 1 \times 10^{-5}$; hypermethylation of cg04605617 (chr1: 20,501,558) and cg08042135 (chr1: 20,501,758) in *PLA2G2C* was observed (Spearman correlation=0.53). However, upon adjustment of cg08042135 by cg04605617, the associations with blood ($P=0.14$) and urinary total arsenic ($P=0.24$) no longer persisted. Sensitivity analyses were conducted with arsenic exposure as a natural log-transformed variable and did not appreciably alter the association P values (data not shown), suggesting robust linear associations for reported loci. Additionally, the methylation β values are presented by urinary total arsenic exposure quartiles in Supplementary Material, Figure S2. In general, dose-dependent trends were observed between urinary total arsenic quartiles and DNA methylation levels. Boxplots looked similar for blood arsenic quartiles (data not shown).

Among all 471,932 CpG loci evaluated, 56.6% of the methylation probes were hypermethylated (t-statistic >0) and 43.4% of methylation probes were hypomethylated (t-statistic <0) in relation

to urinary total arsenic concentration. Results were similar in relation to blood arsenic concentration (data not shown). Among the top 35 differentially methylated loci, blood and urinary total arsenic levels were associated with both gene-specific DNA hypermethylation (N=19; 54.3%) as well as gene-specific DNA hypomethylation (N=16; 45.7%), as shown in Table 2 (P=0.78 for enrichment). Furthermore, no statistically significant association of arsenic exposure was observed in relation to global methylation levels across autosomes on a genomic scale. In a global analysis evaluating the association between arsenic exposure and average β values across all available CpG sites, significant global methylation patterns were not observed for blood (P=0.124) or urinary total arsenic (P=0.241) concentrations.

Among the 35 differentially methylated loci with $P < 1 \times 10^{-5}$, 29 loci could be evaluated in the replication sample. Of these, 8 methylation loci were associated with $P < 0.05$, with 5 loci observed to have the same direction of effect as the discovery association (Supplemental Material, Table S1). The strongest replication signal was observed for *IGH* cg13651690 ($P = 5.40 \times 10^{-3}$). The Kolmogorov test indicated that the replication P values were significantly different from a uniform P value distribution ($P = 0.0011$).

Methylation probes containing known single nucleotide polymorphism (SNPs) were not removed from analyses *a priori* since genome-wide SNP data were available for the study sample (Pierce et al. 2012) and could be examined in stratified analyses. As shown in Table 2, among the significantly differentially methylated loci based on Bonferroni criteria, one probe contained a SNP with minor allele frequency ≥ 0.01 . Methylation probe cg04605617 contained SNP rs12139100 for which genotype data were available on 363 of the 400 study participants. The A allele frequency in the study sample was 0.38. In stratified analyses, the association between

blood and urinary total arsenic concentrations in relation to DNA methylation at cg04605617 was observed to be independent of rs12139100 genotype. Among individuals with the GG genotype (n=136), associations persisted in relation to DNA methylation for blood arsenic concentration ($P=3.08\times 10^{-4}$) and urinary total arsenic concentration ($P=6.20\times 10^{-5}$) as well as among individuals with the GA+AA genotype (n=227) for blood arsenic concentration ($P=1.61\times 10^{-7}$) and urinary total arsenic concentration ($P=2.43\times 10^{-7}$).

Correlation between white blood cell DNA methylation with corresponding peripheral blood mononuclear cell gene expression was examined for the top 35 differentially methylated probes among the 400 study participants. Gene expression signals based on RNA transcripts for the corresponding genes containing the differentially methylated loci are summarized in Table 3. Among the 35 differentially methylated loci with $P<1\times 10^{-5}$, we could evaluate corresponding RNA transcript levels from the same gene for 28 methylation loci. Of these, there were 15 methylation loci significantly associated with gene expression based on $P<0.1$.

To evaluate the potential effect of arsenic exposure on white blood cell type proportions, we utilized estimated cell type fractions based on a validated subset of the methylation data, presented by quartiles of arsenic exposure in Table 4. We observed no notable effect of arsenic on cell type proportions, except for moderate estimated percent decreases in CD4+ T cells and NK cells in relation to the highest quartile of urinary total arsenic concentration as well as borderline associations ($P=0.13-0.14$) in relation to the highest quartile of blood arsenic concentration.

Discussion

The findings of this study suggest associations of blood and urinary total arsenic concentrations with gene-specific DNA methylation changes. We identified four novel methylation loci in *PLA2G2C*, *SQSTM1*, *SLC44A4*, and *IGH* that were strongly associated with arsenic exposure, as well as several suggestive associations in other gene regions. Additionally, we show that several of the differentially methylated loci were associated with corresponding gene expression levels in peripheral blood mononuclear cells.

Higher arsenic exposure was associated with increased methylation levels at cg04605617 (chr1: 20,501,558), located in the first exon of *PLA2G2C*. This locus was moderately associated with increased gene expression of *PLA2G2C* (ILMN_3237030 P=0.073). *PLA2G2C* encodes a calcium-dependent phospholipase, which is an enzyme involved in the hydrolysis of phospholipids into free fatty acids and lysophospholipids. These lipid mediators have diverse biological functions relevant for cancer progression including roles in inflammation, cell growth, signaling and death (Dennis et al. 2011; Scott et al. 2010). Notably, phospholipase A2 enzymes have been previously shown to be induced by skin carcinogens (e.g. phorbol ester and ultraviolet B light), which ultimately lead to prostaglandin synthesis via cyclooxygenase-2 (COX2) leading to increased keratinocyte proliferation and skin carcinogenesis (Bowden 2004; Kast et al. 1993). Arsenic is an established skin carcinogen, with evidence from animal studies suggesting overexpression of COX2 associated with arsenic exposure (Ouyang et al. 2007; Tokar et al. 2011; Trouba and Germolec 2004).

Higher arsenic exposure was associated with decreased methylation levels at the cg01225779 locus (chr5: 179,238,473), located in the 5' UTR of *SQSTM1*. *SQSTM1* encodes a protein that

binds ubiquitin and regulates activation of the nuclear factor kappa-B signaling pathway. *SQSTM1* has been implicated in a number of diseases including neurodegenerative diseases, cancer, obesity and insulin resistance (Geetha et al. 2012). *In vitro* studies have shown that arsenic induces the nuclear factor kappa-B pathway and may be a potential mechanism for skin carcinogenesis (Liao et al. 2004; Zuo et al. 2012).

Higher arsenic exposure was associated with decreased methylation levels at the cg06121226 locus (chr4: 72,134,061). It is located in the body of *SLC4A4*, and was strongly associated with increased gene expression of *SLC4A4* (ILMN_2356991 $P=2.173 \times 10^{-4}$ and ILMN_2184556 $P=0.029$). *SLC4A4* encodes a sodium bicarbonate cotransporter involved in the regulation of bicarbonate secretion and absorption as well as intracellular pH. Mutations in this gene have been associated with hypertension (Yang et al. 2012), a well-established health outcome associated with arsenic exposure (Abhyankar et al. 2012).

Higher arsenic exposure was associated with increased methylation levels at the cg13651690 locus (chr14: 106,320,748), located in the body of *IGH*. The immunoglobulin heavy locus includes variable (V), diversity (D), joining (J), and constant (C) segments of immunoglobulins. Translocations in this region have been implicated in lymphoma (Guais et al. 2004). Furthermore, arsenic exposure has been associated with elevated serum immunoglobulins (Islam et al. 2007), which may be involved in skin carcinogenesis (Wiemels et al. 2011). Furthermore, we observed significant replication of this locus in an independent study sample in relation to water arsenic concentration.

The biological implications of these findings must be further explored with regard to their role in various mechanisms of arsenic toxicity and arsenic-related disease outcomes. Chronic exposure

to arsenic in drinking water has been associated with a multitude of health effects including increased risks of cancer, cardiovascular disease, peripheral neuropathy, and respiratory diseases (Brouwer et al. 1992; Chen et al. 1992; Milton and Rahman 2002; Navas-Acien et al. 2005) as well as a possible association with diabetes (Argos et al. 2013; Navas-Acien et al. 2006). Future studies should be designed to evaluate phenotype-specific methylation patterns in arsenic-exposed populations. Additionally, the magnitude of differential methylation at each locus associated with arsenic exposure was small, although similar in size to estimated effects previously reported for arsenic and other environmental exposures (Joubert et al. 2012; Koestler et al. 2013a). The biological implications of relatively small changes in DNA methylation need to be elucidated further.

Among the 7 epigenome-wide studies in relation to arsenic-related traits published to date (Bailey et al. 2013; Kile et al. 2014; Koestler et al. 2013a; Liu et al. 2014; Seow et al. 2014; Smeester et al. 2011; Yang et al. 2014), all but 2 of the studies utilized the Illumina platform for measurement of DNA methylation (Bailey et al. 2013; Smeester et al. 2011). We did not observe any overlap between our top 35 differentially methylation loci and the top signals previously reported for other arsenic-related traits among studies utilizing the Illumina methylation array; although none of these prior studies identified statistically differentially methylated loci based on a Bonferroni threshold ($P < 1 \times 10^{-7}$). We additionally conducted a lookup of the top reported CpG signals from the previously published studies in our dataset (Supplemental Material, Table S2). Nominally significant associations in our dataset were observed for *AGAP2* cg11511175 and *RHBDF1* cg03333116 previously reported in relation to arsenical skin lesion status (Seow et al. 2014); *ELL* cg22489759 and *SNRNP200* cg00088989 previously reported in relation to toenail

arsenic concentration (Liu et al. 2014); *RIN2* cg03512414, *SLC12A6* cg11293029, *CBFA2T3* cg09051215, and *CCDC73* cg01717164 previously reported in relation to in utero arsenic exposure (Koestler et al. 2013a). Furthermore, we evaluated methylation of genes previously reported based on candidate promoter methylation studies (Banerjee et al. 2013; Chanda et al. 2013; Gribble et al. 2014; Hossain et al. 2012; Intarasunanont et al. 2012) using a lookup approach in our dataset for *DAPK1*, *CDKN2A (P16)*, *GMD5*, *C10orf32/AS3MT*, *RASSF1*, *PPARG*, *TP53*, and *MLH1* (Supplemental Material, Table S3). Our data provide supporting evidence for differential methylation specifically in the promoter regions of *CDKN2A* (cg03079681), *RASSF1* (cg06117233), *TP53* (cg05479194, cg02855142, cg08119584, and cg01620719), and *MLH1* (cg11291081 and cg05670953) in relation to arsenic exposure (Supplemental Material, Table S3). Given the notable differences in arsenic toxicity constructs, arsenic exposure levels, and participant populations across published studies, future research is needed to further synthesize the existing evidence and elucidate the role of epigenetic mechanisms in relation to arsenic exposure and related-diseases.

There are limitations of this study that we also consider. DNA methylation was measured in total white blood cells, which comprise various leukocyte subtypes known to be associated with differential methylation signatures (Adalsteinsson et al. 2012; Reinius et al. 2012). If arsenic exposure was associated with a substantial shift in leukocyte subtypes, then our analyses of DNA methylation in white blood cells may be confounded due to differences in cell type proportions. Since frozen unfractionated blood was utilized in these experiments, we could not evaluate the association between arsenic and cell type fractions directly. Therefore, we utilized a novel statistical method to infer expected cell type fractions in our study samples based on a validated

subset of methylation markers as a surrogate measure (Houseman et al. 2012); the assumptions of the statistical method have been described elsewhere (Koestler et al. 2013a; Koestler et al. 2013b). Based on these analyses, arsenic did not appear to be strongly associated with cell type shifts, except for associations of the highest quartile of urinary total arsenic concentration with decreased CD4+ T and NK cell fractions. Therefore, we do not believe that the results observed in our analyses can be fully explained through an immunotoxic pathway of arsenic.

Another potential consideration for the findings of this study is that all participants had manifest arsenic skin lesions, which is a proxy for both chronic arsenic exposure and genetic susceptibility to arsenic toxicity. Since genotype is known to influence DNA methylation patterns (Tycko 2010), it is possible that the associations observed in this study may not be generalizable to populations without skin lesions. However, since individuals with arsenical skin lesions are at increased risk of developing arsenic-related cancers and other disease conditions, the results of this study offer valuable insight into potential mechanistic pathways related to arsenic toxicity and carcinogenesis. Another potential limitation is that we did not validate the methylation signals identified with the Illumina platform using additional confirmatory methods. Given previously published studies indicating very good concordance of the Illumina 450K platform with pyrosequencing data (Roessler et al. 2012), we did not pursue validation methods in this study.

The major strengths of this study were the relatively large size of the study sample, the multiple measures of arsenic exposure, the broad exposure range, and the availability of epigenome-wide methylation data as well as genome-wide expression and genetic data from the study sample. Whereas previous studies have demonstrated an association between arsenic exposure and DNA

methylation, we were additionally enabled to evaluate potentially functional gene regulation associated with the differentially methylated loci.

Conclusions

In conclusion, we found that arsenic exposure was associated with differential gene-specific white blood cell DNA methylation at several novel loci. We also observed functional evidence of gene deregulation that corresponded with differential methylation at a subset of these loci. However, the clinical implications of these findings in arsenic-exposed populations require further investigation.

References

- Abhyankar LN, Jones MR, Guallar E, Navas-Acien A. 2012. Arsenic exposure and hypertension: A systematic review. *Environ Health Perspect* 120:494-500.
- Adalsteinsson BT, Gudnason H, Aspelund T, Harris TB, Launer LJ, Eiriksdottir G, et al. 2012. Heterogeneity in white blood cells has potential to confound DNA methylation measurements. *PloS One* 7:e46705.
- Argos M, Rahman M, Parvez F, Dignam J, Islam T, Quasem I, et al. 2013. Baseline comorbidities in a skin cancer prevention trial in bangladesh. *Eur J Clin Invest* 43:579-588.
- Bailey KA, Wu MC, Ward WO, Smeester L, Rager JE, Garcia-Vargas G, et al. 2013. Arsenic and the epigenome: Interindividual differences in arsenic metabolism related to distinct patterns of DNA methylation. *J Biochem Mol Toxicol* 27:106-115.
- Banerjee N, Paul S, Sau TJ, Das JK, Bandyopadhyay A, Banerjee S, et al. 2013. Epigenetic modifications of *dapk* and *p16* genes contribute to arsenic-induced skin lesions and nondermatological health effects. *Toxicol Sci* 135:300-308.
- Bowden GT. 2004. Prevention of non-melanoma skin cancer by targeting ultraviolet-b-light signalling. *Nat Rev Cancer* 4:23-35.
- Brouwer OF, Onkenhout W, Edelbroek PM, de Kom JF, de Wolff FA, Peters AC. 1992. Increased neurotoxicity of arsenic in methylenetetrahydrofolate reductase deficiency. *Clin Neurol Neurosurg* 94:307-310.
- Chanda S, Dasgupta UB, Guhamazumder D, Gupta M, Chaudhuri U, Lahiri S, et al. 2006. DNA hypermethylation of promoter of gene *p53* and *p16* in arsenic-exposed people with and without malignancy. *Toxicol Sci* 89:431-437.
- Chanda S, Dasgupta UB, Mazumder DG, Saha J, Gupta B. 2013. Human *gmds* gene fragment hypermethylation in chronic high level of arsenic exposure with and without arsenic induced cancer. *Springerplus* 2:557.
- Chen CJ, Chen CW, Wu MM, Kuo TL. 1992. Cancer potential in liver, lung, bladder and kidney due to ingested inorganic arsenic in drinking water. *Br J Cancer* 66:888-892.
- Chen WT, Hung WC, Kang WY, Huang YC, Chai CY. 2007. Urothelial carcinomas arising in arsenic-contaminated areas are associated with hypermethylation of the gene promoter of the death-associated protein kinase. *Histopathology* 51:785-792.

- Dennis EA, Cao J, Hsu YH, Magrioti V, Kokotos G. 2011. Phospholipase a2 enzymes: Physical structure, biological function, disease implication, chemical inhibition, and therapeutic intervention. *Chem Rev* 111:6130-6185.
- Engstrom KS, Hossain MB, Lauss M, Ahmed S, Raqib R, Vahter M, et al. 2013. Efficient arsenic metabolism--the as3mt haplotype is associated with DNA methylation and expression of multiple genes around as3mt. *PloS One* 8:e53732.
- Geetha T, Vishwaprakash N, Sycheva M, Babu JR. 2012. Sequestosome 1/p62: Across diseases. *Biomarkers* 17:99-103.
- Gribble MO, Tang WY, Shang Y, Pollak J, Umans JG, Francesconi KA, et al. 2014. Differential methylation of the arsenic (iii) methyltransferase promoter according to arsenic exposure. *Arch Toxicol* 88:275-282.
- Guais A, Solhonne B, Melaine N, Guellaen G, Bulle F. 2004. Goliath, a ring-h2 mitochondrial protein, regulated by luteinizing hormone/human chorionic gonadotropin in rat leydig cells. *Biol Reprod* 70:204-213.
- Harper KN, Peters BA, Gamble MV. 2013. Batch effects and pathway analysis: Two potential perils in cancer studies involving DNA methylation array analysis. *Cancer Epidemiol Biomarkers Prev* 22:1052-1060.
- Heinegard D, Tiderstrom G. 1973. Determination of serum creatinine by a direct colorimetric method. *Clin Chim Acta* 43:305-310.
- Hossain MB, Vahter M, Concha G, Broberg K. 2012. Environmental arsenic exposure and DNA methylation of the tumor suppressor gene p16 and the DNA repair gene mlh1: Effect of arsenic metabolism and genotype. *Metallomics* 4:1167-1175.
- Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, et al. 2012. DNA methylation arrays as surrogate measures of cell mixture distribution. *BMC Bioinformatics* 13:86.
- Hughes MF. 2006. Biomarkers of exposure: A case study with inorganic arsenic. *Environ Health Perspect* 114:1790-1796.
- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. 2012. Arsenic, metals, fibres, and dusts. *IARC Monogr Eval Carcinog Risks Hum* 100:11-465.

- Intarasunanont P, Navasumrit P, Waraprasit S, Chaisatra K, Suk WA, Mahidol C, et al. 2012. Effects of arsenic exposure on DNA methylation in cord blood samples from newborn babies and in a human lymphoblast cell line. *Environ Health* 11:31.
- Islam LN, Nabi AH, Rahman MM, Zahid MS. 2007. Association of respiratory complications and elevated serum immunoglobulins with drinking water arsenic toxicity in human. *J Environ Sci Health* 42:1807-1814.
- Johnson WE, Li C, Rabinovic A. 2007. Adjusting batch effects in microarray expression data using empirical bayes methods. *Biostatistics* 8:118-127.
- Joubert BR, Haberg SE, Nilsen RM, Wang X, Vollset SE, Murphy SK, et al. 2012. 450k epigenome-wide scan identifies differential DNA methylation in newborns related to maternal smoking during pregnancy. *Environ Health Perspect* 120:1425-1431.
- Kast R, Furstenberger G, Marks F. 1993. Phorbol ester tpa- and bradykinin-induced arachidonic acid release from keratinocytes is catalyzed by a cytosolic phospholipase a2 (cpla2). *J Invest Dermatol* 101:567-572.
- Kilaru V, Barfield RT, Schroeder JW, Smith AK, Conneely KN. 2012. Methlab: A graphical user interface package for the analysis of array-based DNA methylation data. *Epigenetics* 7:225-229.
- Kile ML, Baccarelli A, Hoffman E, Tarantini L, Quamruzzaman Q, Rahman M, et al. 2012. Prenatal arsenic exposure and DNA methylation in maternal and umbilical cord blood leukocytes. *Environ Health Perspect* 120:1061-1066.
- Kile ML, Houseman EA, Baccarelli A, Quamruzzaman Q, Rahman M, Mostofa G, et al. 2014. Effect of prenatal arsenic exposure on DNA methylation and leukocyte subpopulations in cord blood. *Epigenetics* 9:774-782.
- Kitchin KT, Conolly R. 2010. Arsenic-induced carcinogenesis--oxidative stress as a possible mode of action and future research needs for more biologically based risk assessment. *Chem Res Toxicol* 23:327-335.
- Koestler DC, Avissar-Whiting M, Houseman EA, Karagas MR, Marsit CJ. 2013a. Differential DNA methylation in umbilical cord blood of infants exposed to low levels of arsenic in utero. *Environ Health Perspect* 121:971-977.

- Koestler DC, Christensen B, Karagas MR, Marsit CJ, Langevin SM, Kelsey KT, et al. 2013b. Blood-based profiles of DNA methylation predict the underlying distribution of cell types: A validation analysis. *Epigenetics* 8:816-826.
- Lambrou A, Baccarelli A, Wright RO, Weisskopf M, Bollati V, Amarasiwardena C, et al. 2012. Arsenic exposure and DNA methylation among elderly men. *Epidemiol* 23:668-676.
- Liao WT, Chang KL, Yu CL, Chen GS, Chang LW, Yu HS. 2004. Arsenic induces human keratinocyte apoptosis by the fas/fas ligand pathway, which correlates with alterations in nuclear factor-kappa b and activator protein-1 activity. *J Invest Dermatol* 122:125-129.
- Liu X, Zheng Y, Zhang W, Zhang X, Lioyd-Jones DM, Baccarelli AA, et al. 2014. Blood methylomics in response to arsenic exposure in a low-exposed us population. *J Expo Sci Environ Epidemiol* 24:145-149.
- Majumdar S, Chanda S, Ganguli B, Mazumder DN, Lahiri S, Dasgupta UB. 2010. Arsenic exposure induces genomic hypermethylation. *Environ Toxicol* 25:315-318.
- Marsit CJ, Karagas MR, Danaee H, Liu M, Andrew A, Schned A, et al. 2006. Carcinogen exposure and gene promoter hypermethylation in bladder cancer. *Carcinogenesis* 27:112-116.
- Milton AH, Rahman M. 2002. Respiratory effects and arsenic contaminated well water in Bangladesh. *Int J Environ Health Res* 12:175-179.
- Navas-Acien A, Sharrett AR, Silbergeld EK, Schwartz BS, Nachman KE, Burke TA, et al. 2005. Arsenic exposure and cardiovascular disease: A systematic review of the epidemiologic evidence. *Am J Epidemiol* 162:1037-1049.
- Navas-Acien A, Silbergeld EK, Streeter RA, Clark JM, Burke TA, Guallar E. 2006. Arsenic exposure and type 2 diabetes: A systematic review of the experimental and epidemiological evidence. *Environ Health Perspect* 114:641-648.
- Nixon DE, Mussmann GV, Eckdahl SJ, Moyer TP. 1991. Total arsenic in urine: Palladium-persulfate vs nickel as a matrix modifier for graphite furnace atomic absorption spectrophotometry. *Clin Chem* 37:1575-1579.
- Ouyang W, Zhang D, Ma Q, Li J, Huang C. 2007. Cyclooxygenase-2 induction by arsenite through the ikkbeta/nfkappab pathway exerts an antiapoptotic effect in mouse epidermal cl41 cells. *Environ Health Perspect* 115:513-518.

- Pierce BL, Kibriya MG, Tong L, Jasmine F, Argos M, Roy S, et al. 2012. Genome-wide association study identifies chromosome 10q24.32 variants associated with arsenic metabolism and toxicity phenotypes in Bangladesh. *PLoS Genet* 8:e1002522.
- Pilsner JR, Liu X, Ahsan H, Ilievski V, Slavkovich V, Levy D, et al. 2007. Genomic methylation of peripheral blood leukocyte DNA: Influences of arsenic and folate in Bangladeshi adults. *Am J Clin Nutr* 86:1179-1186.
- Pilsner JR, Liu X, Ahsan H, Ilievski V, Slavkovich V, Levy D, et al. 2009. Folate deficiency, hyperhomocysteinemia, low urinary creatinine, and hypomethylation of leukocyte DNA are risk factors for arsenic-induced skin lesions. *Environ Health Perspect* 117:254-260.
- Pilsner JR, Hall MN, Liu X, Ilievski V, Slavkovich V, Levy D, et al. 2012. Influence of prenatal arsenic exposure and newborn sex on global methylation of cord blood DNA. *PloS One* 7:e37147.
- Pruszkowski E, Neubauer K, Thomas R. 1998. An overview of clinical applications by inductively coupled plasma mass spectrometry. *Atom Spectrosc* 19:111-115.
- Reichard JF, Puga A. 2010. Effects of arsenic exposure on DNA methylation and epigenetic gene regulation. *Epigenomics* 2:87-104.
- Reinius LE, Acevedo N, Joerink M, Pershagen G, Dahlen SE, Greco D, et al. 2012. Differential DNA methylation in purified human blood cells: Implications for cell lineage and studies on disease susceptibility. *PloS One* 7:e41361.
- Ren X, McHale CM, Skibola CF, Smith AH, Smith MT, Zhang L. 2011. An emerging role for epigenetic dysregulation in arsenic toxicity and carcinogenesis. *Environ Health Perspect* 119:11-19.
- Roessler J, Ammerpohl O, Gutwein J, Hasemeier B, Anwar SL, Kreipe H, et al. 2012. Quantitative cross-validation and content analysis of the 450k DNA methylation array from Illumina, Inc. *BMC Res Notes* 5:210.
- Scott KF, Sajinovic M, Hein J, Nixdorf S, Galettis P, Liauw W, et al. 2010. Emerging roles for phospholipase a2 enzymes in cancer. *Biochimie* 92:601-610.
- Seow WJ, Kile ML, Baccarelli AA, Pan WC, Byun HM, Mostofa G, et al. 2014. Epigenome-wide DNA methylation changes with development of arsenic-induced skin lesions in Bangladesh: a case-control follow-up study. *Environ Mol Mutagen* 55:449-456.

- Smeester L, Rager JE, Bailey KA, Guan X, Smith N, Garcia-Vargas G, et al. 2011. Epigenetic changes in individuals with arsenicosis. *Chem Res Toxicol* 24:165-167.
- Smith AH, Lingas EO, Rahman M. 2000. Contamination of drinking-water by arsenic in Bangladesh: A public health emergency. *Bull World Health Org* 78:1093-1103.
- Stroh A. 1988. Determination of Pb and Cd in whole blood using isotope dilution icp-ms. *Atom Spectrosc* 14:141-143.
- Tajuddin SM, Amaral AF, Fernandez AF, Rodriguez-Rodero S, Rodriguez RM, Moore LE, et al. 2013. Genetic and non-genetic predictors of line-1 methylation in leukocyte DNA. *Environ Health Perspect* 121:650-656.
- Tokar EJ, Diwan BA, Ward JM, Delker DA, Waalkes MP. 2011. Carcinogenic effects of "whole-life" exposure to inorganic arsenic in cd1 mice. *Toxicol Sci* 119:73-83.
- Trouba KJ, Germolec DR. 2004. Micromolar concentrations of sodium arsenite induce cyclooxygenase-2 expression and stimulate p42/44 mitogen-activated protein kinase phosphorylation in normal human epidermal keratinocytes. *Toxicol Sci* 79:248-257.
- Tycko B. 2010. Allele-specific DNA methylation: Beyond imprinting. *Hum Mol Genet* 19:R210-220.
- Van Geen A, Ahsan H, Horneman AH, Dhar RK, Zheng Y, Hussain I, et al. 2002. Promotion of well-switching to mitigate the current arsenic crisis in Bangladesh. *Bull World Health Org* 80:732-737.
- Wiemels JL, Wiencke JK, Li Z, Ramos C, Nelson HH, Karagas MR. 2011. Risk of squamous cell carcinoma of the skin in relation to ige: A nested case-control study. *Cancer Epidemiol Biomarkers Prev* 20:2377-2383.
- Wilhelm CS, Kelsey KT, Butler R, Plaza S, Gagne L, Zens MS, et al. 2010. Implications of line1 methylation for bladder cancer risk in women. *Clin Cancer Res* 16:1682-1689.
- Yang HC, Liang YJ, Chen JW, Chiang KM, Chung CM, Ho HY, et al. 2012. Identification of *igf1*, *slc4a4*, *wwox*, and *sfbmt1* as hypertension susceptibility genes in Han Chinese with a genome-wide gene-based association study. *PloS One* 7:e32907.
- Yang TY, Hsu LI, Chiu AW, Pu YS, Wang SH, Liao YT, et al. 2014. Comparison of genome-wide DNA methylation in urothelial carcinomas of patients with and without arsenic exposure. *Environ Res* 128:57-63.

- Zhang AH, Bin HH, Pan XL, Xi XG. 2007. Analysis of p16 gene mutation, deletion and methylation in patients with arseniasis produced by indoor unventilated-stove coal usage in Guizhou, China. *J Toxicol Environ Health* 70:970-975.
- Zuo Z, Ouyang W, Li J, Costa M, Huang C. 2012. Cyclooxygenase-2 (cox-2) mediates arsenite inhibition of uvb-induced cellular apoptosis in mouse epidermal cl41 cells. *Curr Cancer Drug Targets* 12:607-616.

Table 1. Selected characteristics of study sample.

| Characteristic | N (%)^a |
|--|--------------------------|
| Sex | |
| Male | 212 (53.0) |
| Female | 188 (47.0) |
| Age, years | |
| Mean \pm SD | 43.3 \pm 10.2 |
| 25-35 | 104 (26.0) |
| 36-45 | 130 (32.5) |
| 46-55 | 122 (30.5) |
| 56-65 | 44 (11.0) |
| Blood arsenic concentration, $\mu\text{g/L}$ | |
| Mean \pm SD | 9.3 \pm 11.3 |
| 0.80-2.50 | 110 (27.7) |
| 2.51-4.79 | 88 (22.2) |
| 4.80-11.20 | 100 (25.2) |
| 11.21-81.60 | 99 (24.9) |
| Urinary total arsenic concentration, $\mu\text{g/g}$ | |
| Mean \pm SD | 302 \pm 364.5 |
| 12.0-75.0 | 97 (24.2) |
| 75.1-139.9 | 105 (26.3) |
| 140.0-394.9 | 98 (24.5) |
| 395.0-2250.0 | 100 (25.0) |

SD, standard deviation.

^aAll values are N (%) unless otherwise denoted.

Table 2. Top 35 differentially methylated loci based on $P < 1 \times 10^{-5}$ in relation to blood or urinary total arsenic concentrations, sorted by chromosome.

| Probe | Chr | Position | Gene | Feature category ^a | Median β value | Blood arsenic concentration | | | Urinary total arsenic concentration | | |
|-------------------------|-----|-------------|-----------------|-------------------------------|-------------------------|-----------------------------|-------|------------------------|-------------------------------------|-------|------------------------|
| | | | | | | Coeff. | SE | P value | Coeff. | SE | P value |
| cg02856716 | 1 | 18,993,307 | PAX7 | Body | 0.102 | -0.014 | 0.004 | 2.28×10^{-4} | -0.019 | 0.004 | 6.46×10^{-6} |
| cg04605617 ^b | 1 | 20,501,558 | PLA2G2C | 1st Exon | 0.724 | 0.049 | 0.007 | 1.48×10^{-11} | 0.054 | 0.008 | 3.40×10^{-11} |
| cg08042135 | 1 | 20,501,758 | PLA2G2C | TSS200 | 0.774 | 0.019 | 0.004 | 2.71×10^{-6} | 0.020 | 0.004 | 1.37×10^{-5} |
| cg13223043 | 1 | 26,492,308 | FAM110D ZNF593 | Intergenic | 0.448 | -0.021 | 0.004 | 1.48×10^{-6} | -0.022 | 0.005 | 6.78×10^{-6} |
| cg00857921 | 1 | 92,257,380 | TGFB3 | Body | 0.393 | 0.016 | 0.004 | 4.12×10^{-5} | 0.021 | 0.004 | 1.98×10^{-6} |
| cg19750321 | 1 | 150,808,974 | ARNT | Body | 0.853 | 0.017 | 0.004 | 7.06×10^{-6} | 0.016 | 0.004 | 2.20×10^{-4} |
| cg03984502 | 1 | 151,805,662 | RORC | TSS1500 | 0.816 | 0.020 | 0.004 | 6.44×10^{-6} | 0.015 | 0.005 | 2.34×10^{-3} |
| cg07207669 | 1 | 155,102,389 | EFNA1 | Body | 0.560 | -0.023 | 0.005 | 5.90×10^{-7} | -0.025 | 0.005 | 1.27×10^{-6} |
| cg08438392 | 2 | 27,708,645 | IFT172 | Body | 0.782 | 0.020 | 0.004 | 3.27×10^{-6} | 0.022 | 0.005 | 3.66×10^{-6} |
| cg00522451 | 2 | 113,464,049 | SLC20A1 NT5DC4 | Intergenic | 0.216 | -0.025 | 0.005 | 2.49×10^{-7} | -0.028 | 0.005 | 3.16×10^{-7} |
| cg00281776 ^c | 2 | 209,224,226 | PIKFYVE PTH2R | Intergenic | 0.817 | -0.038 | 0.008 | 1.22×10^{-6} | -0.042 | 0.009 | 1.50×10^{-6} |
| cg25881170 | 3 | 107,810,508 | CD47 | TSS1500 | 0.473 | 0.019 | 0.004 | 3.28×10^{-6} | 0.022 | 0.005 | 1.14×10^{-6} |
| cg24262469 | 3 | 156,391,694 | TIPARP-AS1 | Body | 0.371 | 0.022 | 0.005 | 8.48×10^{-6} | 0.022 | 0.005 | 4.98×10^{-5} |
| cg06121226 | 4 | 72,134,061 | SLC4A4 | Body | 0.463 | -0.054 | 0.009 | 7.63×10^{-9} | -0.059 | 0.010 | 1.16×10^{-8} |
| cg01225779 | 5 | 179,238,473 | SQSTM1 | 5'UTR | 0.562 | -0.042 | 0.007 | 6.00×10^{-9} | -0.048 | 0.008 | 2.37×10^{-9} |
| cg19937878 | 6 | 13,296,150 | TBC1D7 | Body | 0.820 | 0.028 | 0.007 | 1.91×10^{-5} | 0.035 | 0.007 | 1.59×10^{-6} |
| cg11230112 ^d | 6 | 158,460,980 | SYNJ2 | Body | 0.872 | -0.029 | 0.006 | 6.60×10^{-7} | -0.029 | 0.007 | 1.54×10^{-5} |
| cg22813794 | 7 | 75,677,469 | MDH2 | TSS200; 5'UTR | 0.039 | -0.059 | 0.014 | 5.49×10^{-5} | -0.077 | 0.016 | 1.54×10^{-6} |
| cg15310871 | 8 | 20,077,937 | ATP6V1B2 | 3'UTR | 0.825 | -0.039 | 0.008 | 3.61×10^{-7} | -0.042 | 0.008 | 1.09×10^{-6} |
| cg22977892 | 8 | 25,907,769 | PPP2R2A | Body | 0.254 | 0.021 | 0.005 | 7.70×10^{-6} | 0.020 | 0.005 | 1.20×10^{-4} |
| cg02268561 | 10 | 15,212,066 | NMT2 | TSS1500 | 0.118 | -0.026 | 0.005 | 6.23×10^{-7} | -0.026 | 0.006 | 7.74×10^{-6} |
| cg15108641 | 10 | 99,263,321 | UBTD1 | Body | 0.365 | 0.033 | 0.006 | 3.70×10^{-7} | 0.030 | 0.007 | 3.11×10^{-5} |
| cg07545081 | 10 | 126,308,381 | FAM53B | 3'UTR | 0.685 | 0.021 | 0.005 | 3.42×10^{-6} | 0.026 | 0.005 | 3.68×10^{-7} |
| cg02742555 | 11 | 15,200,693 | INSC | Body | 0.163 | -0.028 | 0.005 | 2.22×10^{-7} | -0.026 | 0.006 | 2.74×10^{-5} |
| cg03348792 | 12 | 53,075,482 | KRT1 | TSS1500 | 0.309 | 0.019 | 0.004 | 6.62×10^{-6} | 0.019 | 0.005 | 8.91×10^{-5} |
| cg06383241 | 12 | 116,997,023 | MAP1LC3B2 | TSS200 | 0.083 | -0.022 | 0.005 | 2.29×10^{-6} | -0.022 | 0.005 | 2.23×10^{-5} |
| cg11582226 | 13 | 77,587,297 | FBXL3 | Body | 0.756 | 0.021 | 0.004 | 3.40×10^{-6} | 0.019 | 0.005 | 1.98×10^{-4} |
| cg13651690 | 14 | 106,320,748 | IGH | Body | 0.950 | 0.035 | 0.006 | 1.24×10^{-7} | 0.039 | 0.007 | 9.16×10^{-8} |
| cg05018460 | 15 | 80,688,079 | AB240015 ARNT2 | Intergenic | 0.557 | 0.047 | 0.010 | 1.56×10^{-6} | 0.046 | 0.011 | 2.09×10^{-5} |
| cg04352288 ^e | 16 | 87,958,408 | CA5A | Body | 0.872 | 0.047 | 0.011 | 9.20×10^{-6} | 0.056 | 0.012 | 2.60×10^{-6} |
| cg17892169 | 17 | 7,452,644 | TNFSF12-TNFSF13 | Body | 0.341 | 0.024 | 0.005 | 8.06×10^{-6} | 0.023 | 0.006 | 2.01×10^{-4} |
| cg08285388 | 17 | 27,230,177 | DHRS13 | TSS200 | 0.090 | -0.014 | 0.004 | 1.37×10^{-3} | -0.022 | 0.005 | 3.29×10^{-6} |
| cg13480898 | 19 | 10,195,915 | C19orf66 | TSS1500 | 0.680 | -0.023 | 0.005 | 5.35×10^{-6} | -0.026 | 0.005 | 1.63×10^{-6} |
| cg06381803 | 19 | 46,119,476 | EML2 | Body | 0.400 | -0.042 | 0.010 | 5.72×10^{-5} | -0.051 | 0.011 | 8.95×10^{-6} |
| cg26390598 | 21 | 41,032,397 | B3GALT5 | 5'UTR | 0.340 | 0.050 | 0.011 | 9.03×10^{-6} | 0.063 | 0.012 | 5.49×10^{-7} |

Probes containing SNPs with a MAF >0.01 reported. All SNPs >10b from query site. “||” indicates an intergenic region. TSS200, 200 bases from transcription start site (TSS). TSS1500, 1500 bases from TSS. UTR, untranslated region.

^aFeature categories indicate the gene feature category of the methylation locus. ^brs12139100. ^crs139141387. ^drs73795212. ^ers113904153.

Table 3. Top 35 differentially methylated loci based on $P < 1 \times 10^{-5}$ in relation to peripheral blood mononuclear cell gene expression, sorted by chromosome.

| Probe | Chr | Position | Gene | Feature category ^a | Probe | Probe coordinates | P value | Direction ^b |
|------------|-----|-------------|------------------------|-------------------------------|--|--|--|------------------------|
| cg02856716 | 1 | 18,993,307 | <i>PAX7</i> | Body | ILMN_1835658 ILMN_1761061 | 19,074,941-19,074,990 19,062,405-19,062,454 | 0.574 0.818 | + + |
| cg04605617 | 1 | 20,501,558 | <i>PLA2G2C</i> | 1st Exon | ILMN_3237030 ILMN_1656867 | 20,501,573-20,501,622 20,490,548-20,490,597 | 0.073 0.324 | + - |
| cg08042135 | 1 | 20,501,758 | <i>PLA2G2C</i> | TSS200 | ILMN_1656867 ILMN_3237030 | 20,490,548-20,490,597 20,501,573-20,501,622 | 0.742 0.758 | - + |
| cg13223043 | 1 | 26,492,308 | <i>FAM110D ZNF593</i> | Intergenic | NA | | | |
| cg00857921 | 1 | 92,257,380 | <i>TGFB3</i> | Body | ILMN_1784287 | 92,148,163-92,148,212 | 1.620×10^{-8} | - |
| cg19750321 | 1 | 150,808,974 | <i>ARNT</i> | Body | ILMN_1762582 ILMN_2347314 | 150,782,288-150,782,337 150,782,989-150,783,038 | 0.351 0.552 | + - |
| cg03984502 | 1 | 151,805,662 | <i>RORC</i> | TSS1500 | ILMN_1771126 ILMN_1734366 ILMN_1651792 ILMN_2275399 | 151,778,991-151,779,040 151,778,901-151,778,950 151,804,215-151,804,264 151,798,406-151,798,455 | 3.001×10^{-6} 4.770×10^{-6} 0.247 0.992 | - - + - |
| cg07207669 | 1 | 155,102,389 | <i>EFNA1</i> | Body | ILMN_2371055 ILMN_2371053 | 155,107,215-155,107,264 155,106,804-155,106,853 | 0.140 0.272 | + - |
| cg08438392 | 2 | 27,708,645 | <i>IFT172</i> | Body | ILMN_1784178 | 27,668,277-27,668,621 | 0.428 | - |
| cg00522451 | 2 | 113,464,049 | <i>SLC20A1 NT5DC4</i> | Intergenic | NA | | | |
| cg00281776 | 2 | 209,224,226 | <i>PIKFYVE PTH2R</i> | Intergenic | NA | | | |
| cg25881170 | 3 | 107,810,508 | <i>CD47</i> | TSS1500 | ILMN_2356991 ILMN_1771333 | 107,762,694-107,762,743 107,762,387-107,762,436 | 0.044 0.060 | + + |
| cg24262469 | 3 | 156,391,694 | <i>TIPARP-AS1</i> | Body | ILMN_3239662 | 157,875,849-157,875,878 | 0.223 | + |
| cg06121226 | 4 | 72,134,061 | <i>SLC4A4</i> | Body | ILMN_1734897 ILMN_2184556 | 72,437,058-72,437,107 72,437,334-72,437,383 | 2.173×10^{-4} 0.029 | + + |
| cg01225779 | 5 | 179,238,473 | <i>SQSTM1</i> | 5'UTR | NA | | | |
| cg19937878 | 6 | 13,296,150 | <i>TBC1D7</i> | Body | ILMN_1661622 | 13,305,354-13,305,403 | 0.567 | - |
| cg11230112 | 6 | 158,460,980 | <i>SYNJ2</i> | Body | ILMN_2215119 | 158,438,833-158,438,882 | 0.752 | + |
| cg22813794 | 7 | 75,677,469 | <i>MDH2</i> | TSS200; 5'UTR | ILMN_2079004 | 75,695,852-75,695,901 | 0.867 | + |
| cg15310871 | 8 | 20,077,937 | <i>ATP6V1B2</i> | 3'UTR | ILMN_1787705 | 20,078,945-20,078,994 | 0.004 | - |
| cg22977892 | 8 | 25,907,769 | <i>PPP2R2A</i> | Body | NA | | | |
| cg02268561 | 10 | 15,212,066 | <i>NMT2</i> | TSS1500 | ILMN_2062620 | 15,147,945-15,147,994 | 0.030 | + |
| cg15108641 | 10 | 99,263,321 | <i>UBTD1</i> | Body | ILMN_1794914 | 99,330,592-99,330,641 | 2.416×10^{-6} | - |
| cg07545081 | 10 | 126,308,381 | <i>FAM53B</i> | 3'UTR | ILMN_2053490 ILMN_1704571 | 126,308,101-126,308,150 126,308,260-126,308,309 | 0.280 0.834 | - - |

| Probe | Chr | Position | Gene | Feature category ^a | Probe | Probe coordinates | P value | Direction ^b |
|------------|-----|-------------|------------------------|-------------------------------|--------------|-------------------------|---------|------------------------|
| cg02742555 | 11 | 15,200,693 | <i>INSC</i> | Body | ILMN_1756070 | 15,262,019-15,262,068 | 0.343 | + |
| | | | | | ILMN_2340643 | 15,267,549-15,267,598 | 0.983 | + |
| cg03348792 | 12 | 53,075,482 | <i>KRT1</i> | TSS1500 | ILMN_1735712 | 53,068,865-53,068,914 | 0.298 | + |
| cg06383241 | 12 | 116,997,023 | <i>MAP1LC3B2</i> | TSS200 | ILMN_3247613 | 115,481,580-115,481,621 | 0.005 | – |
| cg11582226 | 13 | 77,587,297 | <i>FBXL3</i> | Body | ILMN_2071405 | 77,579,718-77,579,767 | 0.299 | + |
| cg13651690 | 14 | 106,320,748 | <i>IGH</i> | Body | NA | | | |
| cg05018460 | 15 | 80,688,079 | <i>AB240015 ARNT2</i> | Intergenic | NA | | | |
| cg04352288 | 16 | 87,958,408 | <i>CA5A</i> | Body | ILMN_1731292 | 87,921,734-87,921,783 | 0.004 | + |
| cg17892169 | 17 | 7,452,644 | <i>TNFSF12-TNFSF13</i> | Body | ILMN_1683700 | 7,460,967-7,461,016 | 0.022 | – |
| | | | | | ILMN_2399190 | 7,462,480-7,462,529 | 0.031 | + |
| | | | | | ILMN_1784264 | 7,464,109-7,464,339 | 0.150 | + |
| | | | | | ILMN_1670188 | 7,452,806-7,453,453 | 0.335 | – |
| | | | | | ILMN_1680003 | 7,457,089-7,457,138 | 0.819 | + |
| cg08285388 | 17 | 27,230,177 | <i>DHRS13</i> | TSS200 | ILMN_1790781 | 27,224,816-27,224,865 | 0.366 | – |
| cg13480898 | 19 | 10,195,915 | <i>C19orf66</i> | TSS1500 | ILMN_1750400 | 10,203,832-10,203,881 | 0.004 | + |
| cg06381803 | 19 | 46,119,476 | <i>EML2</i> | Body | ILMN_3240541 | 50,804,842-50,804,886 | 0.028 | – |
| cg26390598 | 21 | 41,032,397 | <i>B3GALT5</i> | 5'UTR | ILMN_1698756 | 39,954,012-39,954,061 | 0.059 | – |
| | | | | | ILMN_2378654 | 39,956,034-39,956,083 | 0.206 | – |
| | | | | | ILMN_1800713 | 39,956,436-39,956,485 | 0.376 | – |

NA indicates that a corresponding RNA expression transcript was not available. “||” indicates an intergenic region. TSS200, 200 bases from transcription start site (TSS). TSS1500, 1500 bases from TSS. UTR, untranslated region.

^aFeature categories indicate the gene feature category of the methylation locus. ^bA “–” indicates an inverse association whereas a “+” indicates a positive association between methylation and gene expression levels

Table 4. Estimated change in leukocyte cell type proportions by blood and urinary total arsenic concentrations.

| Exposure | CD8+ T cells | CD4+ T cells | NK cells | B cells | Monocytes | Granulocytes |
|---|--------------------|-----------------------------------|-----------------------------------|---------------------|---------------------|---------------------|
| Blood arsenic concentration (µg/L) | | | | | | |
| 0.80-2.50 | Ref | Ref | Ref | Ref | Ref | Ref |
| 2.51-4.79 | 0.39 (-1.09, 1.87) | -0.57 (-1.59, 0.44) | 0.05 (-0.06, 0.15) | -0.46 (-1.18, 0.26) | 0.04 (-0.47, 0.55) | 0.55 (-1.16, 2.26) |
| 4.80-11.20 | 1.02 (-0.43, 2.47) | -0.30 (-1.29, 0.70) | 0.02 (-0.08, 0.12) | -0.39 (-1.09, 0.32) | -0.05 (-0.54, 0.45) | -0.31 (-1.99, 1.37) |
| 11.21-81.60 | 0.47 (-0.97, 1.90) | -0.75 (-1.74, 0.24) | -0.08 (-0.18, 0.02) | -0.41 (-1.11, 0.29) | 0.15 (-0.34, 0.64) | 0.61 (-1.05, 2.28) |
| Urinary total arsenic concentration (µg/g) | | | | | | |
| 12.0-75.0 | Ref | Ref | Ref | Ref | Ref | Ref |
| 75.1-139.9 | 0.37 (-1.08, 1.81) | -0.72 (-1.71, 0.28) | -0.08 (-0.18, 0.02) | -0.31 (-1.01, 0.39) | 0.25 (-0.24, 0.75) | 0.49 (-1.18, 2.16) |
| 140.0-394.9 | 1.21 (-0.27, 2.69) | -0.47 (-1.50, 0.55) | -0.02 (-0.13, 0.08) | -0.35 (-1.07, 0.37) | 0.20 (-0.31, 0.70) | -0.56 (-2.28, 1.15) |
| 395.0-2250.0 | 0.99 (-0.47, 2.45) | -1.18 (-2.19, -0.17) ^a | -0.12 (-0.22, -0.01) ^a | -0.68 (-1.39, 0.03) | 0.18 (-0.32, 0.68) | 0.81 (-0.88, 2.50) |

^aP<0.05.

Figure Legends

Figure 1. Manhattan plot for epigenome-wide association study results for blood arsenic concentration. The horizontal red line corresponds to significance threshold $P=1\times 10^{-7}$.

Figure 2. Manhattan plot for epigenome-wide association study results for urinary total arsenic concentration. The horizontal red line corresponds to significance threshold $P=1\times 10^{-7}$.

Figure 1.




Figure 2.

